

RESEARCH ARTICLE


Exercise induced plasma volume expansion lowers cardiovascular strain during 15-km cycling time-trial in acute normobaric hypoxia

Felipe Gorini Pereira^{1,2*}, Andrew M. Greenfield¹, Matthew Kuennen³, Trevor L. Gillum¹

1 Department of Kinesiology, California Baptist University, Riverside, CA, United States of America,

2 Department of Kinesiology, Indiana University, Bloomington, IN, United States of America, **3** Department of Exercise Science, High Point University, High Point, NC, United States of America

* fgpereir@iu.edu



Abstract

The purpose of our study was to assess the influence of a single high-intensity interval exercise (HIIE) bout in normoxia on plasma volume (PV) and consequent cycling performance in normobaric hypoxia (0.15 F_iO₂, simulating ~2,500 m). Eight males (VO_{2peak}: 48.8 ± 3.4 mL/kg/min, 24.0 ± 1.6 years) completed a hypoxic 15 km cycling time trial (TT), followed by a crossover intervention of either HIIE (8x4 min cycling bouts at 85% of VO_{2peak}) or CON (matched kJ production from HIIE at 50% of VO_{2peak}). 48 hours post intervention, an identical TT was performed. Cardiovascular parameters were measured via impedance cardiography during each TT. Changes in PV was measured 24 and 48 hours post HIIE and CON. HIIE increased PV at 24 (4.1 ± 3.9%, *P* = 0.031) and 48 (6.7 ± 1.7, *P* = 0.006) hours post, while no difference was observed following the CON (1.3 ± 1.1% and 0.3 ± 2.8%). The higher PV led to an increased stroke volume (*P* = 0.03) and cardiac output (*P* = 0.02) during the hypoxic TT, while heart rate was not changed (*P* = 0.49). We observed no changes in time to completion (−0.63 ± 0.57 min, *P* = 0.054) and power output (7.37 ± 7.98 W, *P* = 0.078) between TTs. In the absence of environmental stress, a single bout of HIIE was an effective strategy to increase PV and reduce the cardiovascular strain during a cycling TT at moderate simulated altitude but did not impact hypoxic exercise performance.

Trial registration: Clinical Trials ID: [NCT05800808](https://clinicaltrials.gov/ct2/show/study/NCT05800808)

OPEN ACCESS

Citation: Gorini Pereira F, Greenfield AM, Kuennen M, Gillum TL (2024) Exercise induced plasma volume expansion lowers cardiovascular strain during 15-km cycling time-trial in acute normobaric hypoxia. PLoS ONE 19(2): e0297553. <https://doi.org/10.1371/journal.pone.0297553>

Editor: Emiliano Cè, Università degli Studi di Milano: Università degli Studi di Milano, ITALY

Received: June 7, 2023

Accepted: January 9, 2024

Published: February 2, 2024

Copyright: © 2024 Gorini Pereira et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data utilized for this study can be freely accessed within the Indiana University Public Repository Database using the following link: https://datacore.iu.edu/concern/data_sets/8p58pd562?locale=en.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Arterial oxygen saturation and content is reduced during dynamic exercise in hypoxic conditions, necessitating increased blood flow in order to meet the augmented energetic demand of the locomotor muscles [1, 2]. Moreover, respiratory muscle fatigue is exacerbated in this environment, as previous work suggests that the work of breathing during intense exercise is augmented during hypoxia exposure [3], congruently forcing respiratory muscles to perform increased work output with limited blood flow. These detrimental factors lead to an exuberant increase in cardiovascular demand, necessitating an increased heart rate (HR) to offset the lower oxygen saturation, in an attempt to maintain adequate cardiac output (\dot{Q}) in light of an

Abbreviations: BL, Blood Lactate; CON, Control; Hb, Hemoglobin; Hct, Hematocrit; HIIE, High Intensity Interval Exercise; HR, Heart Rate; PV, Plasma Volume; Q, Cardiac Output; RPE, Rating of Perceived Exertion; SpO₂, Arterial Oxygen Saturation; SV, Stroke Volume; TT, Time Trial; USG, Urine Specific Gravity; VO_{2peak}, Maximal Oxygen Uptake.

accelerated rate of peripheral fatigue [4]. For this reason, unacclimated individuals experience decreased work capacity during acute altitude exposure [5–7]. Many turn to conventional altitude acclimatization strategies that incorporate intermittent altitude exposures. Best practice guidelines currently recommend 14–19 days for endurance performance [8, 9], but this most likely varies based upon the hours associated with a given level of hypoxia [10]. However, acclimatization is often impractical due to the amount of time required and a lack of access to proper training facilities. As a result, military personnel, wildland firefighters, collegiate, and recreational athletes often perform physical activities at moderate altitude in an unacclimated state.

Interest in novel strategies that could help to mitigate altitude-associated declines in exercise capacity has grown over the years. Elevations in plasma volume (PV) enhance end diastolic ventricular filling and thus \dot{Q} , contributing to reductions in cardiovascular strain at a given exercise intensity [11]. Indeed, acute PV expansion has been shown to reduce cardiovascular strain during sea level exercise [12, 13]. It is also thought to improve prolonged endurance exercise performance, which under normal conditions is characterized by significant reductions in PV [14]. Studies have utilized intravenous infusions of pharmacological agents (i.e. dextran, Gelofusine, and glycerol) to increase PV, which has been shown to improve maximal oxygen uptake ($\dot{V}O_{2peak}$) and prolong exercise-time to exhaustion [15, 16]. Importantly, despite noticeable reductions in hemoglobin (Hb) and hematocrit (Hct) concentrations, these improvements in performance are well established. It has been suggested that perhaps the elevations in SV and \dot{Q} that accompany expansion of PV may offset the unfavorable effects of a reduction in Hb concentration, thereby allowing muscle oxygen delivery to be preserved during high intensity exercise [12, 17]. Given these findings, an expansion of PV may be a beneficial strategy to partially mitigate the deleterious effects of hypoxia on the cardiovascular system.

Rather than focusing on PV expansion via pharmacological manipulation that may have limited application outside laboratory environments, we question if high intensity interval exercise (HIIE) may be an adequate stimulus to promote hypervolemia in a recreationally active cohort. Green et al. demonstrated a 12% increase in PV following three days of intermittent supramaximal exercise bouts [18]. Furthermore, Gillen et al. described a PV increase of ~10% 24 hours following a single session of eight, 4-min intervals at 85% of $\dot{V}O_{2peak}$ [19]. These findings suggest that HIIE could be a practical and time-effective strategy to induce PV expansion, as compared to the utilization of pharmacological agents and/or heat-exposure strategies. However, to our knowledge, no reports have attempted to explore if HIIE could be utilized as a rapid means to expand PV and if these effects might confer benefits during subsequent exercise performance in moderate hypoxia. To this end, we have characterized the PV expansion response from a single HIIE protocol [19] and ascertained the impact of said hypervolemic response on time-trial (TT) performance in normobaric hypoxia (F_iO₂: 0.15, ~2,500 m). We hypothesized that a single HIIE session would increase PV and attenuate cardiovascular strain during exercise in hypoxia, as evidenced by reductions in HR and elevations in SV and \dot{Q} . Furthermore, we hypothesized that these changes would contribute to a reduced time-to-completion in a 15 km, self-paced cycling TT.

Methods and materials

Eight recreationally active males were recruited to participate in this study (Table 1). Sample size was determined with G*Power 3.1.9.4 [20] using Δ PV (evidenced by variations in Hct & Hb) from a recreationally active pool of subjects 24 hours following a single running session of

Table 1. Subject demographics (N = 8).

	Mean	SD
Age (years)	24.0	1.8
Weight (kg)	80.2	9.7
Height (cm)	182.3	7.1
BMI	23.6	3.5
Body Fat (%)	12.3	4.2
$\dot{V}O_{2peak}$ (ml/kg/min ⁻¹)	48.8	3.4
Peak Power (W)	395	48.6

Data are represented as mean \pm SD.

<https://doi.org/10.1371/journal.pone.0297553.t001>

eight, 4-min intervals at 85% of $\dot{V}O_{2peak}$ ($\Delta PV = 10.1 \pm 0.9\%$) [19]. From those data, it was determined that an α -level of $P \leq 0.05$, seven participants would provide an 80% probability (i.e., $1 - \beta$) of detecting a statistically significant change in PV following the prescribed HIIE bout.

Participants were screened for environmental exposure to heat and hypoxia and were not frequent hot bath or sauna users in the three months prior to data collection. Experimental trials were performed in late fall through spring months (Riverside, CA) where weather was not favorable for natural heat acclimatization. Further, the elevation of Riverside, CA is 252 meters above sea level. Participants were screened for cardiovascular and metabolic disease using the American College of Sports Medicine risk stratification guidelines and a Physical Activity Readiness Questionnaire [21]. Participants were not taking any form of medication or supplements throughout the duration of the study. Study staff verbally informed each participant of research procedures, and each gave written informed consent as declared in the 1964 Declaration of Helsinki. All protocols were approved by the Institutional Review Board of California Baptist University (IRB 072-1819-EXP) and the study was registered on clinicaltrials.gov (NCT05800808). All participants were assigned an identification number, which was then used when recording and analyzing data. A code list containing the participant's name and identification numbers were stored separately in a locked filing cabinet and was only available to investigators.

A crossover design was utilized to assess the effects of exercise-induced PV expansion on cycling performance in hypoxia. All participants completed HIIE and a control intervention (CON) in a counterbalanced order. HIIE consisted of 8 x 4 min bouts of cycling exercise at 85% of $\dot{V}O_{2peak}$ with 4 min rest between intervals (19). CON consisted of cycling at 50% $\dot{V}O_{2peak}$ and the overall workload was matched to the HIIE intervention in kilojoules (kJ) produced, resulting in similar workout output for both training conditions (HIIE = 546 ± 84.1 kJ; CON = 547 ± 82.3 kJ). The duration of the CON bout was extended to properly match HIIE kJ production (HIIE: 60 ± 0.0 min; CON: 81 ± 7.2 min). The HIIE and CON training bouts were performed in a thermoneutral and normoxic environment (20°C, 35% RH, $F_iO_2 = 0.2093$). Two-15 km, self-paced cycling TTs were performed before and after each training intervention (Fig 1). The first TT occurred 5 days before the training intervention (HIIE or CON) and the second TT occurred 48 hours post intervention. Interventions were separated by 14 days to ensure sufficient washout of any training effect (Fig 1) as data have suggested retention of expanded PV may last for 7–14 days [22]. During washout, participants were instructed to continue their normal exercise routine.




Fig 1. Study schematic and design. Study participants were randomized into HIIE or CON intervention groups. On visits 1 and 4, both groups completed hypoxic TT-1 and TT-2 for their respective intervention. Following washout and group switch, on visits 5 and 8, both groups completed hypoxic TT-1 and TT-2 for their remaining intervention. *Denotes visits where blood draws occurred.

<https://doi.org/10.1371/journal.pone.0297553.g001>

Preliminary testing

During preliminary testing, body composition was assessed via a 3-site skinfold (chest, abdomen, thigh) measurement. Each site was measured twice in a rotational order and the mean values were summed and used to estimate body density and fat using a standardized regression equation [23]. Aerobic capacity was measured through indirect calorimetry (Parvo Medic, Sandy UT). Participants performed incremental exercise on a cycle ergometer (Monark LC7 TT, Monark, Sweden), where the exercise workload increased 20 Watts per minute following a starting workload of 50 Watts, and pedal cadence was maintained at or above 80 revolutions per minute. For the test to be considered maximum, three of the following $\dot{V}O_{2peak}$ criteria were met: 1) $\dot{V}O_2$ plateau (less than 0.250 L/min), despite increase in workload, 2) rating of perceived exertion (RPE) ≥ 17 , 3) respiratory exchange ratio > 1.10 , and 4) HR ≥ 10 bpm of age-predicted max. Maximal workload was recorded and used to determine exercise workload for HIIE and CON training. After resting for ~15-minutes, participants performed a familiarization 15 km TT in room temperature normoxia (20 C, 35% RH, F_{iO_2} : 0.20). This was done to accustom participants to the equipment and reduce the likelihood of learning effects causing data bias.

Hypoxic TTs

The first hypoxic TT within each condition served as a baseline measure of performance for each participant (Fig 1). Participants abstained from strenuous exercise and alcohol 24 hours prior to each TT. Participants were required to maintain a food log 36 hours prior to the first TT, and then were requested to replicate it for subsequent visits. Upon arrival to the laboratory, participants voided their bladder and provided a urine sample for hydration assessment. Urine specific gravity (USG) was analyzed using a handheld refractometer (ATAGO USA, Bellevue, WA) and adequate hydration was determined as a specified cutoff of USG < 1.020 [24]. Following 20 min of seated rest, a 3 mL blood sample was drawn from the antecubital vein. Three microcuvette tubes were prepared and centrifuged (Bio Lion, Manassas, VA) for later hematological analysis of Hct. The remainder of the blood was immediately dispensed into an EDTA coated vacutainer and analyzed for Hb concentration via photospectrometry (UNICO, Princeton, NJ).

Nitrogen dilution was used to create a hypoxic environment inside an environmental chamber (Cascade Solutions, El Cajon, CA), which was monitored through two independent O_2 sensors (BioSpherix P360) set to maintain F_{iO_2} levels of 0.15 (simulating ~ 2,500 m of elevation).

Participants wore standard exercise attire which included cycling shorts and a t-shirt. Participants were then instrumented with an impedance cardiography apparatus (PhysioFlow, Nanatec Biomedical, France) to measure hemodynamic variables (HR, \dot{Q} , SV). The anatomical

sites required for impedance measurement were cleansed with isopropyl alcohol and shaved (if needed) to ensure proper contact and conductance. Prior to exercise, resting blood pressure was recorded after 20 mins of seated rest. Resting blood pressure was recorded to ensure proper calibration for impedance cardiography measurement. Following instrumentation, participants entered the environmental chamber (20°C, 35% RH) and performed a 10 min warmup by cycling at a power output corresponding to 50% of their individual $\dot{V}O_{2\text{peak}}$.

Upon completion of the warmup, participants performed a 15 km TT as quickly as possible. Throughout the TT, distance covered was made available to participants, but time, speed, cadence, and power were blinded. All participants were instructed to complete the TT as quickly as possible and received verbal encouragement. Blood lactate (Nova Biomedical, Waltham, MA) was measured before (BL_{Rest}) and immediately after (BL_{peak}) each TT from a finger capillary whole blood sample. RPE, which ranged from 6 (no exertion) to 20 (maximal exertion), was also measured via the Borg scale [25] and SpO_2 was measured at the index finger using a standard pulse oximeter (Nellcor Bedside, Nellcor Inc. Hayward, CA).

PV expansion analysis

Blood samples were collected following 20 min seated rest prior to exercise for each TT and testing condition (HIIE and CON). A total of 8 blood samples (4 per intervention) were collected throughout the study as depicted in Fig 1. Hb and Hct were measured to determine changes in PV, as described by Dill and Costill [26] using the following equation:

$$\Delta PV(\%) = \frac{Hb_{\text{pre}} X (1 - Hct_{\text{post}})}{Hb_{\text{post}} X (1 - Hct_{\text{pre}})}$$

Using the above equation, we calculated ΔPV (%) for three distinct time-points: Pre-training (i.e., prior to HIIE/CON), 24 hours post training, and 48 hours post training (i.e., just prior to TT-2).

Statistical analysis

Data were analyzed using GraphPad Prism version 9.0.0 (Prism, San Diego, CA) statistical software. Normality of dependent variables were determined using a Shapiro–Wilk test and homogeneity of variance using Levene’s test. An alpha level of 0.05 was set a priori. Changes in PV were analyzed using a two-factor (intervention x time) repeated measures ANOVA. Averaged hemodynamic parameters (SV (mL/beat), HR (bpm), and \dot{Q} (L/min)), as well as perceptual (RPE_{final}) and physiological (SpO_2 (%), BL_{Rest} (mmol/L), BL_{peak} (mmol/L), and USG) responses during the TT were analyzed using a two-factor (intervention x trial) repeated measure ANOVA. We performed the Holm-Sidak post-hoc procedure to determine where differences existed if the ANOVA revealed a significant interaction or main effect. Mean differences of TT performance variables (time to completion (min) and average power output (W)) were analyzed using two-tailed Wilcoxon matched-pairs, signed rank tests. All data values are reported as mean \pm standard deviation (SD) or mean difference \pm SD. Cohen’s d was calculated to determine effect size with a value of 0.2, 0.5, 0.8 representing small, medium, and large effects; respectively [27].

Results

Plasma volume expansion

Changes in PV throughout the study are shown in Fig 2. There was a significant interaction for PV ($P = 0.017$), which differed between HIIE and CON at 24 hours ($P = 0.031$; $d = 0.49$)




Fig 2. Changes in PV ($n = 8$) 24 and 48 hours following each intervention (HIIE vs. CON). Bars represent mean values for each intervention overlaid by individual values. * Denotes significant differences between interventions at each time point ($P < 0.05$). Data were analyzed using a two-factor (intervention \times trial) repeated measures ANOVA. When a significant main effect or interaction was identified, we performed the Holm-Sidak post-hoc test to determine where differences existed. $n = 8$ for all comparisons.

<https://doi.org/10.1371/journal.pone.0297553.g002>

and 48 hours ($P = 0.006$; $d = 0.71$). No differences for ΔPV were observed between HIIE and CON for pre-training ($P = 0.128$).

There was a significant interaction for Hct ($P = 0.002$), which differed between HIIE and CON at the 48 hours post timepoint ($P = 0.030$; $d = 0.51$; Fig 3A). There were no observed effects of time ($P = 0.212$) or intervention ($P = 0.882$) for Hb (Fig 3B).

Hypoxic cycling time-trial

Averaged hemodynamic parameters for individual TTs are denoted on Table 2. There was a main effect of trial for HR ($P = 0.008$; $d = 0.3$), but no intervention ($P = 0.491$) or interaction ($P = 0.844$) effects were observed. There was an interaction effect for SV ($P = 0.032$; $d = 0.4$), which differed between HIIE TT-1 and TT-2 ($P = 0.029$, $d = 0.4$) and HIIE TT-2 and CON TT-2 ($P = 0.007$; $d = 0.5$). There was an intervention effect for \dot{Q} ($P = 0.023$; $d = 0.3$), which differed between HIIE TT-1 and TT-2 ($P = 0.047$; $d = 0.4$) and HIIE TT-2 and CON TT-2 ($P = 0.027$; $d = 0.7$). RPE_{final} , SpO_2 , BL_{Rest} , BL_{peak} , and USG were not different between HIIE or CON and there were no differences between participant responses at TT-1 and TT-2 (all $P > 0.05$) (Table 2).

Considering all subjects ($n = 8$), Wilcoxon matched-pairs signed rank test revealed no differences in change in time to completion of TT (-0.63 ± 0.57 min, $P = 0.054$; $d = 0.4$; Fig 4A),




Fig 3. A. Hematocrit values ($n = 8$) for all timepoints in which blood sampling occurred. B. Hemoglobin values ($n = 8$) for all timepoints in which blood sampling occurred. Data are expressed as means \pm SD. * Denotes significant differences between interventions at each time point ($P < 0.05$). Data were analyzed using a two-factor (intervention \times trial) repeated measures ANOVA. When a significant main effect or interaction was identified, we performed the Holm-Sidak post-hoc test to determine where differences existed. $n = 8$ for all comparisons.

<https://doi.org/10.1371/journal.pone.0297553.g003>

as well as change in averaged self-selected power output (7.37 ± 7.98 W, $P = 0.078$; $d = 0.7$) for TTs between interventions (Fig 4B).

Discussion

Our main findings are that HIIE increased PV at 24 and 48 hours post exercise (Fig 2), as compared to CON, even though the interventions were matched for total workload. Elevated PV improved hemodynamic responses during TT hypoxic exercise, as evidenced by the significant increases in SV and \dot{Q} without noticeable changes in HR (Table 2). Despite this, there were no differences in time to completion and power output during the TT (Fig 4). These data suggest that HIIE-induced PV expansion lowered cardiovascular strain, and may be an effective strategy to those seeking to mitigate decrements in exercise capacity in moderate hypoxia, without prior acclimation.

Table 2. Hemodynamic, perceptual, and physiological responses to 15 km hypoxic TT.

	TT-1		TT-2	
	HIIE	CON	HIIE	CON
HR (bpm)	165 ± 7.8	167 ± 9.6	$164 \pm 6.6^*$	165 ± 11.2
SV (mL)	125 ± 35	122 ± 36	$133 \pm 34^\dagger$	122 ± 36
Q (L/min)	20.5 ± 5.0	20.2 ± 5.4	$21.9 \pm 4.8^{**}$	20.1 ± 5.2
RPE _{final}	17.0 ± 1.7	17.5 ± 1.9	17.0 ± 1.0	16.6 ± 1.2
SpO ₂ (%)	87.3 ± 1.9	86.3 ± 2.4	86.6 ± 1.4	86.7 ± 1.9
BL _{Rest} (mmol/L)	2.0 ± 0.6	1.8 ± 0.4	2.1 ± 0.7	2.4 ± 0.7
BL _{Peak} (mmol/L)	13.8 ± 2.9	13.5 ± 2.6	14.5 ± 3.9	14.9 ± 2.8
USG	1.010 ± 0.006	1.007 ± 0.004	1.009 ± 0.006	1.008 ± 0.003

Averaged ($n = 8$) hemodynamic parameters for each hypoxic TT. Data are presented as mean \pm SD. USG reflects pre-exercise measurement. *Trial effect ($P < 0.05$).

**Intervention effect ($P < 0.05$).

† Interaction (intervention \times trial) ($P < 0.05$). Data were analyzed using a two-factor (intervention \times trial) repeated measures ANOVA. When a significant main effect or interaction was identified, we performed the Holm-Sidak post-hoc test to determine where differences existed. $n = 8$ for all comparisons.

<https://doi.org/10.1371/journal.pone.0297553.t002>




Fig 4. A. Changes in time to completion of 15-km TT between TT-1 and TT-2; B. Changes in average power output during 15-km TT between TT-1 and TT-2. Bars represent mean differences between TTs for each intervention overlaid by individual values. Data were analyzed using two-tailed Wilcoxon matched-pairs, signed rank tests. $n = 8$ for all comparisons.

<https://doi.org/10.1371/journal.pone.0297553.g004>

HIIE and PV expansion

Exercise training leads to an increase of blood volume, which appears to be primarily accounted for by PV expansion [28]. Cross-sectional analysis of literature indicates PV can be increased within 24 hours following exercise and achieve ~10% increase from pre-exercise levels within 1 to 4 days [29]. The mechanism of hypervolemia associated with acute exercise-training is speculated to be congruently regulated by fluid retention (via renin-angiotensin-aldosterone system) [30], and increases in plasma albumin content [31], as these proteins account for 60–80% of the colloid osmotic pressure in the intravascular space [32].

Our investigation utilized a single HIIE bout, as we sought to isolate intensity-derived PV expansion exclusively developed from our independently matched interventions. To ensure that no changes in PV were due to exercise in hypoxia, PV expansion was measured between TT-1 and HIIE or CON and no differences were revealed (HIIE = $-1.02 \pm 1.94\%$, CON = $0.60 \pm 2.09\%$; $P = 0.128$). Indeed, these data demonstrate that a significant increase in PV was observed 24 (4.05% Δ PV) and 48 (6.87% Δ PV) hours following the HIIE intervention only, further validating existing work documenting the rapid induction of hypervolemia with high-intensity exercise [19, 32, 33]. Accordingly, the degree of PV variation observed in our findings closely resembles those found in the literature surrounding blood volume changes, as Nagashima et al. induced a 6.4% increase in PV 22 hours following a similar upright exercise bout, whilst Gillen's group reported a 10% expansion of PV following a similar treadmill high-intensity exercise bout [19, 32]. Furthermore, the absence of an environmental stressor in the current endeavor points to exercise intensity's pivotal role as a signal for PV expansion, as previously suggested by others [31, 34]. Our findings further corroborate these outcomes, as demonstrated by the lack of Δ PV following the total workload matched CON intervention (-1.20% Δ PV 24 hours post CON; -0.20% Δ PV 48 hours post CON). As a consequence of the augmentation in PV following the HIIE intervention, we observed decreases in Hb and Hct concentrations, both at 24 (Hb = -5.00% ; Hct = -2.91%) and 48 (Hb = -5.57% ; Hct = -4.67%) hours post intervention.

Hemodynamics

The observed PV expansion altered the cardiovascular response to TT performance in acute hypoxia. Improved SV and \dot{Q} following HIT-induced hypervolemia are expected to be a

product of increased left ventricular end-diastolic volume [35], speculated to arise due to enhanced central venous pressure [36]. There are data to support our findings of reduced cardiovascular strain (increased \dot{Q} whilst maintaining exercising HR) following different dosages of known PV expanders (Dextran or Pentaspan) [37, 38], aerobic training [39], and a single bout HIIE protocol resembling our study [40]. These improvements in hemodynamic variables were observed in different testing protocols at various exercising intensities, including at rest [40], during 90 minutes of moderate intensity exercise (62% of $\dot{V}O_{2peak}$) [38], and at different time points throughout a 2-hour low intensity (46% of $\dot{V}O_{2peak}$) cycling bout [37]. Our data are congruent with these studies, but we extend these findings into a hypoxic environment during TT performance. Interestingly, this increase in \dot{Q} is not a consistent finding in literature, as two previous studies have found no effect of artificial PV expansion on \dot{Q} and HR during submaximal exercise, despite observing transient increases in exercise SV [12, 41].

Alongside this reduction in cardiovascular strain, we observed decreases in Hb (24 hours post HIIE = -5.00%; 48 hours post HIIE = -5.57%) and Hct (24 hours post HIIE = -2.91%; 48 hours post HIIE = -4.67%) concentrations (Fig 3). This is an important finding that likely corroborates previous reports suggesting that increases in SV and \dot{Q} , induced via acute PV expansion, are capable of offsetting reductions in Hb and Hct concentrations, such that muscle oxygen delivery is preserved during high intensity exercise [12, 17]. As such, this study highlights that intensity-mediated PV expansion is rapid, which may expedite performance changes when compared to traditional altitude acclimatization protocols that often require ~2 weeks of intermittent hypoxia exposure [42].

TT performance

Various groups regularly travel to moderate altitude (2,000–3,000 m) for competition, recreation, or occupational tasks. The negative effects of hypoxia at this level of altitude are apparent, rapid, and well documented: a 7% reduction in aerobic performance [43], 12% loss in $\dot{V}O_{2peak}$ [44], and an increase in submaximal HR [45]. Thus, there is a need to better understand countermeasures to exercise in acute hypoxia.

Various investigations have shown significant improvements in exercise performance attributed to an acute expansion of PV, making it a relevant adaptation for recreational endurance athletes to consider prior to competition [15, 16, 46, 47]. These data suggest that an enhancement in PV may be of benefit for those seeking incremental improvement of athletic performance at altitude. We observed only a trend in TT performance (time to completion: $P = 0.054$; $d = 0.4$; power output: $P = 0.078$; $d = 0.7$) despite robust increases in PV 24 and 48 hours post interval exercise. While the p values did not cross the *a priori* threshold for significance, the medium to large effect size suggest an improvement in cycling performance was realized. Though our findings differ from previously mentioned reports, our study was not sufficiently powered to directly investigate improvements in exercise performance. Further, data should be interpreted with caution since our participants were recreationally active and thus not accustomed to time trial performance. Our design incorporated a familiarization session prior to the first TT to remove the learning effect of pacing, but the inexperience of participants could have impacted our TT results.

Perspectives

Given the wide variety of endurance sport competition offerings at moderate to high altitudes, questions regarding timing of arrival at altitude are continuously asked by professional and recreational athletes alike. Particularly, one arrival strategy commonly used by national soccer

teams [48] termed “fly in, fly out”, proposes that arrival at altitude should be as close as logistically possible to the event start time, in order to minimize the acute reductions in PV associated with altitude exposure [49]. Previous investigations have observed ~5% reductions in PV within 14 hours of simulated 16% F_iO₂ [50], and > 10% reductions in PV within the first 24 hours at 4,000m of elevation [51, 52]. Reductions in PV are beneficial in increasing the relative concentration of Hb per liter of blood to preserve arterial oxygen content. However, this reduction concomitantly serves to elevate cardiovascular strain via reductions in SV, which can negatively impact exercise performance at altitude [53]. Nevertheless, while the reduction in PV is well-documented upon exposure to > 3,000 m, the minimal altitude at which these reductions take place remain unclear. Reports suggest that exposure to lower altitudes (defined as < 2,500 m) seem to have a minimal effect on PV [45, 54, 55]. In such cases, our findings suggest that an expansion of PV may be a beneficial strategy to partially mitigate the deleterious effects of hypoxia on the cardiovascular system, particularly when exercise will be initiated promptly upon ascent.

Limitations

Few people are afforded the resources to properly acclimate to hypoxia. Given that we sought to evaluate the effectiveness of strategies employed at sea level in better preparing individuals for the enhanced metabolic demands of exercise at altitude, there are inherited limitations associated with this design. The results of this investigation are limited initially by the relatively small cohort ($n = 8$) and exclusive use of a male population. The choice to exclude females from the study was deliberate as it is speculated that the significant fluctuation in ovarian hormones throughout the menstrual cycle can impact blood volume following dynamic exercise [56]. These variations could confound re-creation of our results within a similarly trained female population. Also, the participants could not be blinded to the intervention, as the awareness of intervals was impossible to overcome. Thus, some participants may have expected to perform better after intervals compared to after control. Furthermore, the use of impedance cardiography to assess hemodynamic variables needs to be acknowledged. Previous work has highlighted potential limitations in the use of this technique [57], specifically regarding appropriate placement of electrodes and artifacts from neighboring organs. Despite these concerns, this device and its validity during rest and exercise testing have been previously published [58]. Additionally, placement of electrodes, calibration of device, as well as extraction of the collected data were performed in accordance with the manufacturer’s instructions. Finally, our participants were recreationally active and thus their PV response to HIIE may differ compared to highly trained athletes, who have already expanded their plasma through habitual aerobic training.

Conclusion

A single cycling bout of HIIE is an effective strategy to induce hypervolemia within 24- and 48-hours following exercise. In the present study, this expansion in PV served to increase stroke volume and cardiac output, while lowering HR. The utilization of high intensity intervals prior to athletic endeavors may be an alluring strategy to increase cardiac output during exercise in hypoxia.

Acknowledgments

The authors would like to extend our gratitude to our dedicated study participants.

Author Contributions

Conceptualization: Felipe Gorini Pereira, Trevor L. Gillum.

Data curation: Felipe Gorini Pereira, Andrew M. Greenfield.

Formal analysis: Felipe Gorini Pereira, Andrew M. Greenfield, Trevor L. Gillum.

Investigation: Felipe Gorini Pereira.

Methodology: Felipe Gorini Pereira.

Project administration: Trevor L. Gillum.

Software: Felipe Gorini Pereira.

Supervision: Trevor L. Gillum.

Validation: Felipe Gorini Pereira.

Writing – original draft: Felipe Gorini Pereira, Andrew M. Greenfield, Trevor L. Gillum.

Writing – review & editing: Felipe Gorini Pereira, Andrew M. Greenfield, Matthew Kuennen, Trevor L. Gillum.

References

1. Alexander JK, Hartley LH, Modelski M, Grover RF. Reduction of stroke volume during exercise in man following ascent to 3,100 m altitude. *J Appl Physiol*. 1967; 23(6):849–58. <https://doi.org/10.1152/jappl.1967.23.6.849> PMID: 6065055
2. Fulco CS, Rock PB, Cymerman A. Maximal and submaximal exercise performance at altitude. *Aviat Space Environ Med*. 1998; 69(8):793–801. PMID: 9715971
3. Cibella F, Cuttitta G, Romano S, Grassi B, Bonsignore G, Milic-Emili J. Respiratory energetics during exercise at high altitude. *J Appl Physiol* (1985). 1999; 86(6):1785–92. <https://doi.org/10.1152/jappl.1999.86.6.1785> PMID: 10368338
4. Goodall S, Twomey R, Amann M. Acute and chronic hypoxia: implications for cerebral function and exercise tolerance. *Fatigue*. 2014; 2(2):73–92. <https://doi.org/10.1080/21641846.2014.909963> PMID: 25593787
5. Beidleman BA, Muza SR, Fulco CS, Cymerman A, Sawka MN, Lewis SF, et al. Seven intermittent exposures to altitude improves exercise performance at 4300 m. *Med Sci Sports Exerc*. 2008; 40(1):141–8. <https://doi.org/10.1249/mss.0b013e31815a519b> PMID: 18091011
6. Chapman RF, Stager JM, Tanner DA, Stray-Gundersen J, Levine BD. Impairment of 3000-m run time at altitude is influenced by arterial oxyhemoglobin saturation. *Med Sci Sports Exerc*. 2011; 43(9):1649–56. <https://doi.org/10.1249/MSS.0b013e318211bf45> PMID: 21311361
7. Fulco CS, Muza SR, Beidleman BA, Demes R, Staab JE, Jones JE, et al. Effect of repeated normobaric hypoxia exposures during sleep on acute mountain sickness, exercise performance, and sleep during exposure to terrestrial altitude. *Am J Physiol Regul Integr Comp Physiol*. 2011; 300(2):R428–36. <https://doi.org/10.1152/ajpregu.00633.2010> PMID: 21123763
8. Chapman RF, Karlsen T, Ge RL, Stray-Gundersen J, Levine BD. Living altitude influences endurance exercise performance change over time at altitude. *J Appl Physiol* (1985). 2016; 120(10):1151–8. <https://doi.org/10.1152/jappphysiol.00909.2015> PMID: 26968028
9. Schuler B, Thomsen JJ, Gassmann M, Lundby C. Timing the arrival at 2340 m altitude for aerobic performance. *Scand J Med Sci Sports*. 2007; 17(5):588–94. <https://doi.org/10.1111/j.1600-0838.2006.00611.x> PMID: 17316377
10. Rasmussen P, Siebenmann C, Díaz V, Lundby C. Red cell volume expansion at altitude: a meta-analysis and Monte Carlo simulation. *Med Sci Sports Exerc*. 2013; 45(9):1767–72. <https://doi.org/10.1249/MSS.0b013e31829047e5> PMID: 23502972
11. Fortney SM, Wenger CB, Bove JR, Nadel ER. Effect of blood volume on forearm venous and cardiac stroke volume during exercise. *J Appl Physiol Respir Environ Exerc Physiol*. 1983; 55(3):884–90. <https://doi.org/10.1152/jappl.1983.55.3.884> PMID: 6629925
12. Hopper MK, Coggan AR, Coyle EF. Exercise stroke volume relative to plasma-volume expansion. *J Appl Physiol* (1985). 1988; 64(1):404–8. <https://doi.org/10.1152/jappl.1988.64.1.404> PMID: 2451658

13. Kanstrup IL, Ekblom B. Acute hypervolemia, cardiac performance, and aerobic power during exercise. *J Appl Physiol Respir Environ Exerc Physiol*. 1982; 52(5):1186–91. <https://doi.org/10.1152/jappl.1982.52.5.1186> PMID: 7096143
14. Greenleaf JE, Looft-Wilson R, Wisherd JL, Jackson CG, Fung PP, Ertl AC, et al. Hypervolemia in men from fluid ingestion at rest and during exercise. *Aviat Space Environ Med*. 1998; 69(4):374–86. PMID: 9561285
15. Berger NJ, Campbell IT, Wilkerson DP, Jones AM. Influence of acute plasma volume expansion on VO₂ kinetics, VO₂ peak, and performance during high-intensity cycle exercise. *J Appl Physiol* (1985). 2006; 101(3):707–14. <https://doi.org/10.1152/japplphysiol.00154.2006> PMID: 16690793
16. Coyle EF, Hopper MK, Coggan AR. Maximal oxygen uptake relative to plasma volume expansion. *Int J Sports Med*. 1990; 11(2):116–9. <https://doi.org/10.1055/s-2007-1024774> PMID: 1692570
17. Gledhill N, Warburton D, Jamnik V. Haemoglobin, blood volume, cardiac function, and aerobic power. *Can J Appl Physiol*. 1999; 24(1):54–65. <https://doi.org/10.1139/h99-006> PMID: 9916181
18. Green HJ, Hughson RL, Thomson JA, Sharratt MT. Supramaximal exercise after training-induced hypervolemia. I. Gas exchange and acid-base balance. *J Appl Physiol* (1985). 1987; 62(5):1944–53. <https://doi.org/10.1152/jappl.1987.62.5.1944> PMID: 3597267
19. Gillen CM, Lee R, Mack GW, Tomaselli CM, Nishiyasu T, Nadel ER. Plasma volume expansion in humans after a single intense exercise protocol. *J Appl Physiol* (1985). 1991; 71(5):1914–20. <https://doi.org/10.1152/jappl.1991.71.5.1914> PMID: 1761491
20. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007; 39(2):175–91. <https://doi.org/10.3758/bf03193146> PMID: 17695343
21. American College of Sports M. ACSM's guidelines for exercise testing and prescription: 11th edition. Philadelphia: Lippincott Williams & Wilkins, [2000] ©2000; 2022.
22. Patterson MJ, Stocks JM, Taylor NA. Sustained and generalized extracellular fluid expansion following heat acclimation. *J Physiol*. 2004; 559(Pt 1):327–34. <https://doi.org/10.1113/jphysiol.2004.063289> PMID: 15218070
23. Brozek J, Grande F, Anderson JT, Keys A. Densitometric Analysis of Body Composition: Revision of Some Quantitative Assumptions. *Ann N Y Acad Sci*. 1963; 110:113–40. <https://doi.org/10.1111/j.1749-6632.1963.tb17079.x> PMID: 14062375
24. Armstrong LE, Maresh CM, Castellani JW, Bergeron MF, Kenefick RW, LaGasse KE, et al. Urinary indices of hydration status. *Int J Sport Nutr*. 1994; 4(3):265–79. <https://doi.org/10.1123/ijns.4.3.265> PMID: 7987361
25. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc*. 1982; 14(5):377–81. PMID: 7154893
26. Dill DB, Costill DL. Calculation of percentage changes in volumes of blood, plasma, and red cells in dehydration. *J Appl Physiol*. 1974; 37(2):247–8. <https://doi.org/10.1152/jappl.1974.37.2.247> PMID: 4850854
27. Cohen J. Statistical Power Analysis for The Behavioral Sciences. Hillsdale, NJ: Lawrence Erlbaum Associates; 1988.
28. Oscai LB, Williams BT, Hertig BA. Effect of exercise on blood volume. *J Appl Physiol*. 1968; 24(5):622–4. <https://doi.org/10.1152/jappl.1968.24.5.622> PMID: 5647638
29. Sawka MN, Convertino VA, Eichner ER, Schnieder SM, Young AJ. Blood volume: importance and adaptations to exercise training, environmental stresses, and trauma/sickness. *Med Sci Sports Exerc*. 2000; 32(2):332–48. <https://doi.org/10.1097/00005768-200002000-00012> PMID: 10694114
30. Lavoie JL, Sigmund CD. Minireview: overview of the renin-angiotensin system—an endocrine and paracrine system. *Endocrinology*. 2003; 144(6):2179–83. <https://doi.org/10.1210/en.2003-0150> PMID: 12746271
31. Senay LC, Mitchell D, Wyndham CH. Acclimatization in a hot, humid environment: body fluid adjustments. *J Appl Physiol*. 1976; 40(5):786–96. <https://doi.org/10.1152/jappl.1976.40.5.786> PMID: 931907
32. Nagashima K, Mack GW, Haskell A, Nishiyasu T, Nadel ER. Mechanism for the posture-specific plasma volume increase after a single intense exercise protocol. *J Appl Physiol* (1985). 1999; 86(3):867–73. <https://doi.org/10.1152/jappl.1999.86.3.867> PMID: 10066698
33. Nagashima K, Cline GW, Mack GW, Shulman GI, Nadel ER. Intense exercise stimulates albumin synthesis in the upright posture. *J Appl Physiol* (1985). 2000; 88(1):41–6. <https://doi.org/10.1152/jappl.2000.88.1.41> PMID: 10642360
34. Convertino VA, Brock PJ, Keil LC, Bernauer EM, Greenleaf JE. Exercise training-induced hypervolemia: role of plasma albumin, renin, and vasopressin. *J Appl Physiol Respir Environ Exerc Physiol*. 1980; 48(4):665–9. <https://doi.org/10.1152/jappl.1980.48.4.665> PMID: 6991463

35. Hellsten Y, Nyberg M. Cardiovascular Adaptations to Exercise Training. *Compr Physiol*. 2015; 6(1):1–32. <https://doi.org/10.1002/cphy.c140080> PMID: 26756625
36. Kanstrup IL, Marving J, Hoiland-Carlsen PF. Acute plasma expansion: left ventricular hemodynamics and endocrine function during exercise. *J Appl Physiol* (1985). 1992; 73(5):1791–6. <https://doi.org/10.1152/jappl.1992.73.5.1791> PMID: 1474053
37. Grant SM, Green HJ, Phillips SM, Sutton JR. Effects of acute expansion of plasma volume on cardiovascular and thermal function during prolonged exercise. *Eur J Appl Physiol Occup Physiol*. 1997; 76(4):356–62. <https://doi.org/10.1007/s004210050261> PMID: 9349652
38. Roy BD, Green HJ, Grant SM, Tarnopolsky MA. Acute plasma volume expansion alters cardiovascular but not thermal function during moderate intensity prolonged exercise. *Can J Physiol Pharmacol*. 2000; 78(3):244–50. PMID: 10721816
39. Green HJ, Jones LL, Hughson RL, Painter DC, Farrance BW. Training-induced hypervolemia: lack of an effect on oxygen utilization during exercise. *Med Sci Sports Exerc*. 1987; 19(3):202–6. PMID: 3600233
40. Gillen CM, Nishiyasu T, Langhans G, Weseman C, Mack GW, Nadel ER. Cardiovascular and renal function during exercise-induced blood volume expansion in men. *J Appl Physiol* (1985). 1994; 76(6):2602–10. <https://doi.org/10.1152/jappl.1994.76.6.2602> PMID: 7928889
41. Mier CM, Domenick MA, Turner NS, Wilmore JH. Changes in stroke volume and maximal aerobic capacity with increased blood volume in men women. *J Appl Physiol* (1985). 1996; 80(4):1180–6. <https://doi.org/10.1152/jappl.1996.80.4.1180> PMID: 8926244
42. Millet GP, Roels B, Schmitt L, Woiron X, Richalet JP. Combining hypoxic methods for peak performance. *Sports Med*. 2010; 40(1):1–25. <https://doi.org/10.2165/11317920-000000000-00000> PMID: 20020784
43. Peronnet F, Thibault G, Cousineau DL. A theoretical analysis of the effect of altitude on running performance. *J Appl Physiol* (1985). 1991; 70(1):399–404. <https://doi.org/10.1152/jappl.1991.70.1.399> PMID: 2010398
44. Terrados N. Altitude training and muscular metabolism. *Int J Sports Med*. 1992; 13 Suppl 1:S206–9. <https://doi.org/10.1055/s-2007-1024641> PMID: 1483777
45. Weston AR, Mackenzie G, Tufts MA, Mars M. Optimal time of arrival for performance at moderate altitude (1700 m). *Med Sci Sports Exerc*. 2001; 33(2):298–302. <https://doi.org/10.1097/00005768-200102000-00020> PMID: 11224821
46. Coles MG, Luetkemeier MJ. Sodium-facilitated hypervolemia, endurance performance, and thermoregulation. *Int J Sports Med*. 2005; 26(3):182–7. <https://doi.org/10.1055/s-2004-820989> PMID: 15776333
47. Luetkemeier MJ, Thomas EL. Hypervolemia and cycling time trial performance. *Med Sci Sports Exerc*. 1994; 26(4):503–9. PMID: 7515456
48. Chapman RF, Laymon AS, Levine BD. Timing of arrival and pre-acclimatization strategies for the endurance athlete competing at moderate to high altitudes. *High Alt Med Biol*. 2013; 14(4):319–24. <https://doi.org/10.1089/ham.2013.1022> PMID: 24377334
49. Chapman RF, Stickford JL, Levine BD. Altitude training considerations for the winter sport athlete. *Exp Physiol*. 2010; 95(3):411–21. <https://doi.org/10.1113/expphysiol.2009.050377> PMID: 19837773
50. Foss JL, Constantini K, Mickleborough TD, Chapman RF. Short-term arrival strategies for endurance exercise performance at moderate altitude. *Journal of Applied Physiology*. 2017; 123(5):1258–65. <https://doi.org/10.1152/japplphysiol.00314.2017> PMID: 28818999
51. Beidleman BA, Staab JE, Muza SR, Sawka MN. Quantitative model of hematologic and plasma volume responses after ascent and acclimation to moderate to high altitudes. *Am J Physiol Regul Integr Comp Physiol*. 2017; 312(2):R265–R72. <https://doi.org/10.1152/ajpregu.00225.2016> PMID: 28039191
52. Hannon JP, Shields JL, Harris CW. Effects of altitude acclimatization on blood composition of women. *J Appl Physiol*. 1969; 26(5):540–7. <https://doi.org/10.1152/jappl.1969.26.5.540> PMID: 5781604
53. Siebenmann C, Hug M, Keiser S, Muller A, van Lieshout J, Rasmussen P, et al. Hypovolemia explains the reduced stroke volume at altitude. *Physiol Rep*. 2013; 1(5):e00094. <https://doi.org/10.1002/phy2.94> PMID: 24303166
54. Terzioglu M, Tuna N. Variations in blood volume at 1.85 km altitude. *J Appl Physiol*. 1954; 6(7):417–22. <https://doi.org/10.1152/jappl.1954.6.7.417> PMID: 13117774
55. Brothers MD, Wilber RL, Byrnes WC. Physical fitness and hematological changes during acclimatization to moderate altitude: a retrospective study. *High Alt Med Biol*. 2007; 8(3):213–24. <https://doi.org/10.1089/ham.2007.8308> PMID: 17824822
56. Green DJ, Hopkins ND, Jones H, Thijssen DH, Eijssvogels TM, Yeap BB. Sex differences in vascular endothelial function and health in humans: impacts of exercise. *Exp Physiol*. 2016; 101(2):230–42. <https://doi.org/10.1113/EP085367> PMID: 26663420

57. Sakamoto K, Muto K, Kanai H, Iizuka M. Problems of impedance cardiography. *Med Biol Eng Comput.* 1979; 17(6):697–709. <https://doi.org/10.1007/BF02441549> PMID: 317910
58. Charloux A, Lonsdorfer-Wolf E, Richard R, Lampert E, Oswald-Mammosser M, Mettauer B, et al. A new impedance cardiograph device for the non-invasive evaluation of cardiac output at rest and during exercise: comparison with the "direct" Fick method. *Eur J Appl Physiol.* 2000; 82(4):313–20. <https://doi.org/10.1007/s004210000226> PMID: 10958374